

Hemorrhoids



C A R I S D I A G N O S T I C S H E A L T H I M P R O V E M E N T S E R I E S

What are hemorrhoids? The term hemorrhoids refers to a condition in which the veins around the anus or lower rectum are swollen and inflamed. Hemorrhoids may result from straining to move stool. Other contributing factors include pregnancy, aging, chronic constipation or diarrhea, and anal intercourse. Hemorrhoids are either inside the anus – internal – or under the skin around the anus – external.

What are the symptoms of hemorrhoids? Many anorectal problems, including fissures, fistulae, abscesses, or irritation and itching, also called pruritus ani, have similar symptoms and are incorrectly referred to as hemorrhoids.

Hemorrhoids usually are not dangerous or life threatening. In most cases, hemorrhoidal symptoms will go away within a few days.

Although many people have hemorrhoids, not all experience symptoms. The most common symptom of internal hemorrhoids is bright red blood covering the stool, on toilet paper, or in the toilet bowl. However, an internal hemorrhoid may protrude through the anus outside the body, becoming irritated and painful. This is known as a protruding hemorrhoid.

Symptoms of external hemorrhoids may include painful swelling or a hard lump around the anus that results when a blood clot forms. This condition is known as a thrombosed external hemorrhoid.

In addition, excessive straining, rubbing, or cleaning around the anus may cause irritation with bleeding and/or itching, which may produce a vicious cycle of symptoms. Draining mucus may also cause itching.

How common are hemorrhoids?

Hemorrhoids are common in both men and women. About half of the population has hemorrhoids by age 50.

Hemorrhoids are also common among pregnant women. The

pressure of the fetus on the abdomen, as well as hormonal changes, cause the hemorrhoidal vessels to enlarge. These vessels are also placed under severe pressure during childbirth. For most women, however, hemorrhoids caused by pregnancy are a temporary problem.

How are hemorrhoids diagnosed? A thorough evaluation and proper diagnosis by the doctor is important any time bleeding from the rectum or blood in the stool occurs. Bleeding may also be a symptom of other digestive diseases, including colorectal cancer.


The doctor will examine the anus and rectum to look for swollen blood vessels that indicate hemorrhoids and will also perform a digital rectal exam with a gloved, lubricated finger to feel for abnormalities.

Closer evaluation of the rectum for hemorrhoids requires an exam with an anoscope, a hollow, lighted tube useful for viewing internal hemorrhoids, or a proctoscope, useful for more completely examining the entire rectum.

To rule out other causes of gastrointestinal bleeding, the doctor may examine the rectum and lower colon, or sigmoid, with sigmoidoscopy or the entire colon with

colonoscopy. Sigmoidoscopy and colonoscopy are diagnostic procedures that also involve the use of lighted, flexible tubes inserted through the rectum.

I N T E R N A L H E M O R R H O I D



© 1997-2008 Nucleus Medical Art, Inc. All Rights Reserved

S E E R E V E R S E S I D E

Hemorrhoids



C A R I S D I A G N O S T I C S H E A L T H I M P R O V E M E N T S E R I E S

What is the treatment? Medical treatment of hemorrhoids is aimed initially at relieving symptoms. Measures to reduce symptoms include

- tub baths several times a day in plain, warm water for about 10 minutes
- application of a hemorrhoidal cream or suppository to the affected area for a limited time

Preventing the recurrence of hemorrhoids will require relieving the pressure and straining of constipation. Doctors will often recommend increasing fiber and fluids in the diet. Eating the right amount of fiber and drinking six to eight glasses of fluid – not alcohol – result in softer, bulkier stools. A softer stool makes emptying the bowels easier and lessens the pressure on hemorrhoids caused by straining. Eliminating straining also helps prevent the hemorrhoids from protruding.

Good sources of fiber are fruits, vegetables, and whole grains. In addition, doctors may suggest a bulk stool softener or a fiber supplement such as psyllium (Metamucil) or methylcellulose (Citrucel).

In some cases, hemorrhoids must be treated endoscopically or surgically. These methods are used to shrink and destroy the hemorrhoidal tissue. The doctor will perform the procedure during an office or hospital visit.

A number of methods may be used to remove or reduce the size of internal hemorrhoids. These techniques include:

- **Rubber band ligation.** A rubber band is placed around the base of the hemorrhoid inside the rectum. The band cuts off circulation, and the hemorrhoid withers away within a few days.

- **Sclerotherapy.** A chemical solution is injected around the blood vessel to shrink the hemorrhoid.
- **Infrared coagulation.** A special device is used to burn hemorrhoidal tissue.
- **Hemorrhoidectomy.** Occasionally, extensive or severe internal or external hemorrhoids may require removal by surgery known as hemorrhoidectomy.

How are hemorrhoids prevented? The best way to prevent hemorrhoids is to keep stools soft so they pass easily, thus decreasing pressure and straining, and to empty bowels as soon as possible after the urge occurs. Exercise, including walking, and increased fiber in the diet help reduce constipation and straining by producing stools that are softer and easier to pass.



8400 Esters Boulevard, Suite 190, Irving, Texas 75063
CarisDx.com 800.979.8292

Content provided by the National Digestive Diseases Information Clearinghouse (NDDIC), a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).

This material is intended for patient education and information only. It does not constitute advice, nor should it be taken to suggest or replace professional medical care from your physician. Your treatment options may vary, depending upon medical history and current condition. Only your physician and you can determine your best option. Provided to you as a service by Caris Diagnostics, Inc. ©2008 Caris Diagnostics, Inc. All rights reserved. (01.08)

Caris Diagnostics provides academic-caliber GI and Dermatology anatomic pathology testing services to physicians who treat patients in an ambulatory setting. Its industry-leading team of fellowship and subspecialty trained pathologists provide the highest levels of service to its clinician customers and their patients through its state-of-the-art laboratories, ongoing educational activities, proprietary advanced clinical practice solutions, and rigorous quality assurance programs. In addition, CarisMPI provides advanced analysis of patient samples through prognostic testing services, genomic and proteomic profiling to help guide physicians in the treatment of cancer and other complex diseases. More than 1,500 physicians nationally send their patient's biopsies to Caris Diagnostics at our four laboratories: Irving, Texas; Phoenix, Arizona(2); and Newton, Massachusetts. Additional information is available at www.carisdx.com.

C A R I S D X . C O M

8 0 0 . 9 7 9 . 8 2 9 2